

ON THE GROMOV WIDTH OF HOMOGENEOUS KÄHLER MANIFOLDS

ANDREA LOI AND FABIO ZUDDAS

ABSTRACT. We compute the Gromov width of homogeneous Kähler manifolds with second Betti number equal to one. Our result is based on the recent preprint [4] and on the upper bound of the Gromov width for such manifolds obtained in [6].

1. INTRODUCTION

The Gromov width [3] of a $2n$ -dimensional symplectic manifold (M, ω) is defined as

$$c_G(M, \omega) = \sup\{\pi r^2 \mid B^{2n}(r) \text{ symplectically embeds into } (M, \omega)\}, \quad (1)$$

where

$$B^{2n}(r) = \{(x, y) \in \mathbb{R}^{2n} \mid \sum_{j=1}^n x_j^2 + y_j^2 < r^2\} \quad (2)$$

is the open ball of radius r endowed with the standard symplectic form $\omega_0 = \sum_{j=1}^n dx_j \wedge dy_j$ of \mathbb{R}^{2n} . By Darboux's theorem $c_G(M, \omega)$ is a positive number. Computations and estimates of the Gromov width for various examples have been obtained by several authors (see, e.g. [6] and references therein). The main result of this paper is the following theorem proved in the next section.

2000 Mathematics Subject Classification. 53D05; 53C55; 53D05; 53D45.

Key words and phrases. Gromov width; homogeneous space.

The authors were supported by Prin 2010/11 – Varietà reali e complesse: geometria, topologia e analisi armonica – Italy. The first author was supported by INdAM–GNSAGA - Gruppo Nazionale per le Strutture Algebriche, Geometriche e le loro Applicazioni.

Theorem 1. *Let (M, ω) be a compact homogeneous Kähler manifold such that $b_2(M) = 1$ and ω is normalized so that $\omega(A) = \int_A \omega = \pi$ for the generator $A \in H_2(M, \mathbb{Z})$. Then*

$$c_G(M, \omega) = \pi. \quad (3)$$

The class of manifolds in Theorem 1 includes all Hermitian symmetric space of compact type whose Gromov width has been computed in [5]. We do not know if the assumption on the second Betti number can be dropped.

2. PROOF OF THEOREM 1

The proof of Theorem 1 is mainly based on the lower bound recently obtained by K. Kaveh [4]:

Theorem A *Let X be a smooth complex projective variety embedded in a complex projective space \mathbb{CP}^N . Then*

$$c_G(X, \omega_{FS}) \geq 1, \quad (4)$$

where ω_{FS} denotes the restriction to X of the Fubini–Study Kähler form of \mathbb{CP}^N .

Proof of Theorem 1 The upper bound $c_G(M, \omega) \leq \pi$ is Theorem 1 in [6]. In order to obtain the lower bound $c_G(M, \omega) \geq \pi$, consider the integral Kähler form $\hat{\omega} = \frac{\omega}{\pi}$ on M . Let (L, h) be the holomorphic hermitian line bundle on M such that $\text{Ric}(h) = \hat{\omega}$, where $\text{Ric}(h)$ is the 2-form on M defined by $\text{Ric}(h) = -\frac{i}{2\pi} \partial \bar{\partial} \log h(\sigma, \sigma)$, for a local trivializing holomorphic section σ of L . Let s_0, \dots, s_N be an orthonormal basis for the space of global holomorphic sections $H^0(L)$ of L equipped with the L^2 -scalar product $\langle \cdot, \cdot \rangle$ given by:

$$\langle s, t \rangle = \int_M h(s, t) \frac{\hat{\omega}^n}{n!}, \quad s, t \in H^0(L).$$

Then, it is not hard to see (see, e.g. [1]), due to the homogeneity and simply connectedness of M , that the Kodaira map $k : M \rightarrow \mathbb{CP}^N, x \mapsto [s_0(x) : \dots : s_N(x)]$ is a Kähler immersion, i.e. $k^* \omega_{FS} = \hat{\omega}$. Moreover, in [2, Theorem 3] is proved that such a map is injective, and hence

$(M, \hat{\omega})$ is symplectomorphic to $(k(M), \omega_{FS})$. By Theorem A and by the conformality of the Gromov width one gets

$$c_G(M, \omega) = \pi c_G(M, \hat{\omega}) = \pi c_G(k(M), \omega_{FS}) \geq \pi$$

and the theorem is proved. \square

REFERENCES

- [1] C. Arezzo, A. Loi, *Moment maps, scalar curvature and quantization of Kähler manifolds*, Comm. Math. Phys. 243 (2004), 543-559.
- [2] A. J. Di Scala, H. Hishi, A. Loi, *Kähler immersions of homogeneous Kähler manifolds into complex space forms*, Asian Journal of Mathematics Vol. 16 No. 3 (2012), 479-488.
- [3] M. Gromov, *Pseudoholomorphic curves in symplectic manifolds*, Invent. Math. 82 (1985), no. 2, 307-347, Springer-Verlag (1986).
- [4] K. Kaveh, *Toric degenerations and Gromov width of smooth projective varieties*, <http://arxiv.org/abs/1508.00316>.
- [5] A. Loi, R. Mossa, F. Zuddas, *Symplectic capacities of Hermitian symmetric spaces of compact and non compact type*, to appear in Journal of Symplectic Geometry, <http://arxiv.org/abs/1302.1984>.
- [6] A. Loi, R. Mossa, F. Zuddas, *Some remarks on the Gromov width of homogeneous Hodge manifolds*, Int. J. Geom. Methods Mod. Phys. Vol. 11, No. 2 (2014).

(ANDREA LOI) DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI CAGLIARI (ITALY)

E-mail address: `loi@unica.it`

(FABIO ZUDDAS) DIPARTIMENTO DI MATEMATICA E INFORMATICA, VIA DELLE SCIENZE 206, UDINE (ITALY)

E-mail address: `fabio.zuddas@uniud.it`